arXiv:1110.6889v1 [astro-ph.HE] 31 Oct 2011

A new model for the *Fermi*-LAT Extragalactic Gamma-Ray Background

M. Cavadini
Dipartimento di Scienza e Alta Tecnologia, Universitá dell'Insubria, via Valleggio 11, I-22100 Como, Italy

R. Salvaterra
INAF/IASF-MI, via Bassini 15, 20133 Milano, Italy

F. Haardt
Dipartimento di Scienza e Alta Tecnologia, Universitá dell'Insubria, via Valleggio 11, I-22100 Como, Italy and
INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy

We compute the contribution of blazars (Flat Spectrum Radio Quasars and BL-Lacs) to the *Fermi*-LAT Extragalactic γ-ray Background, assuming a γ-ray luminosity function proportional to the luminosity function of radio galaxies [9] and the Spectral Energy Distribution of blazars given by the so-called *blazar sequence* [11]. We find that blazars account for the ~45 % of the total *Fermi*-LAT EGB. However, at low (<10 GeV) and high (>50 GeV) energies the blazar contribution falls short to account for the observed background light. At E<10 GeV, the Extragalactic γ-ray Background can be fully explained by considering the additional contribution of star-forming galaxies, whereas at very high energies, where the absorption due to Extragalactic Background Light dominates, the observed background calls for an additional unknown component.

I. INTRODUCTION

The Extragalactic γ-ray background (hereafter EGB) represents a fascinating challenge since its first detection by *SAS-2* satellite above 30 MeV [1]. The γ-ray telescope *EGRET* during nine years of operations, measure an isotropic γ-ray emission in the 30 MeV-30 GeV band [2]. The satellite *Fermi*-LAT after one years of observations improves the *EGRET* measurement confirming the existance of an EGB up to 100 GeV [3].

As already found by *EGRET* and confirmed by *Fermi*-LAT [4], blazars are the most common objects in the extragalactic γ-ray sky. Blazars, radio loud AGNs with the jet pointing towards the Earth, are classified as Flat Spectrum Radio Quasars (FSRQs) and BL-Lacs. After three years of observations, *Fermi*-LAT detected approximately 1000 blazars with about the same number of FSRQs and BL-Lacs [4]. It results to be natural computing their contribution to the EGB (e.g. [5–8]). The main ingredients in the EGB calculation are the γ-ray luminosity function (LF) of blazars and their spectral energy distribution (SED). In our model adopt a LF of FRI and FRII radio galaxies [9] and the blazar SEDs using the the so-called *blazar sequence* [10, 11]. We compute the blazar contribution to the *Fermi*-LAT EGB fitting the differential *Fermi*-LAT logN-logS, having as a sole fitting parameter the fraction of radio galaxies beamed towards the Earth, hence appearing as blazars. Beacause the blazar component alone is not able to fully account for the EGB at low energies (<10 GeV), we consider the γ-ray emission of star-forming galaxies. This allows us to fully explain the EGB but the last point at approximately 80 GeV where the EBL absorption strongly dominates. Notice that here we refer to EGB as the sum of the resolved source component and the unresolved and/or truly diffuse emission.

II. BLAZAR CONTRIBUTION

The blazar contribution (in photons s^{-1} cm^{-2} sr^{-1} MeV^{-1}) to the EGB at the observed energy E_0 is

$$I_{\rm blaz}(E_0) \;=\; \frac{1}{4\pi}\int_0^\infty dz\,\frac{dV}{dz}\int_{\log L_\gamma^{\rm min}}^{\log L_\gamma^{\rm max}} d\log L_\gamma\,\frac{d\Phi_\gamma(L_\gamma,z)}{d\log L_\gamma} \times \frac{dn(L_\gamma,z)}{dE}e^{-\tau_{\gamma\gamma}(E_0,z)}, \qquad (1)$$

where $d\Phi_\gamma(L_\gamma,z)/d\log L_\gamma$ is the γ-ray LF and L_γ is νL_ν (in erg/s) at 100 MeV, $dn(L_\gamma,z)/dE$ is the unabsorbed photon flux per unit energy $E=E_0(1+z)$ measured on Earth of a blazar with luminosity L_γ at redshift z, and $\tau_{\gamma\gamma}(E_0,z)$ is the optical depth for $\gamma-\gamma$ absorption. We adopt the Extragalactic Background Light (EBL) model by [13]. In the above equation dV/dz is the comoving cosmological volume. We set $\log L_\gamma^{\rm min}=43.5$ and $\log L_\gamma^{\rm max}=50$.

The number of sources $N(>F_{\rm ph})$ per steradian with photon flux greater than $F_{\rm ph}$ is

$$N(>F_{\rm ph}) = \frac{1}{4\pi}\int_0^\infty dz\,\frac{dV}{dz}\int_{\log L_\gamma^{\rm min}}^{\log L_\gamma^{\rm max}} d\log L_\gamma\,\frac{d\Phi_\gamma(L_\gamma,z)}{d\log L_\gamma}. \qquad (2)$$

The γ-ray LF of blazars is presently uncertain (see e.g. [12]), so that one has to rely on the LFs computed in other bands, e.g., X-rays [6, 14], or radio [7]. We adopt here the radio LF at 151 MHz of FRI and FRII [9], assumed to be the parent populations of blazars:

$$\frac{\Phi_\gamma(L_\gamma,z)}{d\log L_\gamma} = \kappa\;\frac{\Phi_R(L_R,z)}{d\log L_R}, \qquad (3)$$



FIG. 1: The black line represents the fit to the blazar logN- logS measured by Fermi [15] as a sum of BL-Lac component (blue line) and FSRQ component (red line). Black points are the number counts of all the blazars, red points FSRQs and blue points BL-Lacs



FIG. 2: The black points represent the total EGB (resolved and unresolved sources and all the rest), blue line is the FSRQ component, the green line is the BL-Lac component. The red line is the sum of the two components.

where $L_{\rm R}$ is νL_ν at 151 MHz, and the constant κ is the fraction of blazars over all radio galaxies. In order to convert radio into γ-ray luminosity, we rely on the blazar spectral energy distribution (SED). As pointed out in [11], a relation between the radio and/or bolometric luminosity and the energy of the two peaks exists (the so-called *blazar sequence*, see also [10]). We use the SEDs computed by [6, 10].

A. Results

We use the model discussed in the previous section to compute the total contribution of blazars (FSRQs and BL-Lacs) to the *Fermi*-LAT EGB. To this end we fit blazar logN-logS distribution by letting k to vary. We find $\kappa = (3.93 \pm 0.01) \times 10^{-4}$. As shown in Fig. 1, once fitted the total differential logN-logS, our model predict the correct number of BL-Lacs and FSRQs without any change in the radio LF. This fact support the correctness of the initial hypothesis of radio galaxies as parental population of blazars. The number ratio of blazars to radio galaxies κ can be thought as a measure of the beaming factor of the relativistic jet, which in turn is related to the bulk Lorentz factor Γ. From $\kappa \sim 1/2\Gamma^2$ we derive $\Gamma \sim 35$.
Fig. 2 shows the corrisponding contribution of FSRQs and BL-Lac to the total *Fermi*-LAT EGB. The background intensity is found to be $I_{\rm FSRQ} = 4.22 \times 10^{-6}$ph s^{-1}cm^{-2}sr^{-1} and $I_{\rm BL-Lac} = 2.43 \times 10^{-6}$ph s^{-1}cm^{-2}sr^{-1} for FSRQs and BL-Lacs respectively. The total EGB intensity is therefore $I = 6.65 \times 10^{-6}$ph s^{-1}cm^{-2}sr^{-1}, corrisponding to 45% of the one measured by *Fermi*-LAT.
From the slope of the FSRQ and BL-Lac component in Fig. 2 we can see that the main photon index of FSRQs and BL-Lacs resulting from our model are in agreement with the 2LAC [4].
We note that blazars fall short to explain the measured EGB at E<10 GeV and at E>50 GeV. At low energies, the discrepancy can be fully accounted by star-forming galaxies modeled following the recipes by[7], so that only the last point of the *Fermi*-LAT EGB measurement is not reproduced by our model.

III. DISCUSSION

We have computed the overall contribution of blazars to the *Fermi*-LAT EGB.

Our model relies on two assumptions: the radio LF and the blazar SED. In the following we show the difference with the most reliable works on the contribution of blazars to the *Fermi*-LAT EGB.

- We use the radio LF [9]. Fitting the differential logN-logS we obtain just the overall normalization k, without any change on the bright and faint end of LF.



FIG. 3: The star-forming galaxy component (green line), the total blazar component (red line) and the sum (blue line). In black are the *Fermi*-LAT EGB points.

In previous works [6][14] it is assumed a LF in X band with three free parameters: the total normalization, the amount of bolometric radiation emitted in X-ray and the faint end of the X-ray LF. Differently in [7] the radio LF computed by ([16]) is used, changing the faint end to obtain the contribution of FSRQs.

- The second main point of our model is represented by the blazar SED. We assume that the spectra of blazar is fully taken into account by the *blazar sequence* [11]. On the contrary in [7, 8] it is assumed the blazar spectra as a simple and broken power law, respectively. In these works they assume a spectral index distribution peaked on the mean spectral index resulting from observations. The underlying assumption is that the unresolved blazar component has the same index distribution of the resolved component. Using the blazar sequence such assumption is not necessary because the blazar SEDs are fully determined.

The best fit value of the relative number of blazars with respect to radio galaxies can be translated into a bulk Lorentz factor of the relativistic jet $\Gamma \sim 35$, larger than the average value $\Gamma \sim 15$ estimated in [18]. The two values could be reconciled if blazars commonly show secular γ-ray large variability which modulates the 1-year average flux, as recently proposed in [17]. To be consistent with the *Fermi*-LAT points at lower energy, we add the star-forming component to our blazar model. Fitting the *Fermi*-LAT EBG with this two component model (Fig. 3), we constrains the so-called " star formation efficiency of molecular hydrogen" $\xi = 7.0 \times 10^{-10}\mathrm{yr}^{-1}$, which we found well within existing, much looser observational constraints [19]. The point at 80 GeV remains unaccounted by our models calling for an additional (local or low-z) component. An intriguing possibility is represented by annihilation of a dark matter particle. We found, as an example, that a particle of mass $\simeq 0.5$ TeV and cross section $\langle\sigma v\rangle \simeq 5 \times 10^{-26}$ cm^3 s^{-1} can easly accomodate the last data point.

---

[1] Fichtel, C. E., Simpson, G. S., & Thompson, D. J. 1978, baas, 10, 395
[2] Strong, A. W., Moskalenko, I. V., & Reimer, O. 2004, ApJ, 613, 956
[3] Abdo A. A., et al., 2010a, PhRvL, 104, 101101
[4] The Fermi-LAT collaboration, 2011, arXiv, arXiv:1108.1420
[5] Dermer C. D., 2007, ApJ, 659, 958
[6] Inoue, Y., & Totani, T. 2009, ApJ, 702, 523
[7] Stecker F. W., Venters T. M., 2010, arXiv, arXiv:1012.3678
[8] Venters, T. M., & Pavlidou, V. 2011, ApJ 737, 80
[9] Willott C. J., Rawlings S., Blundell K. M., Lacy M., Eales S. A., 2001, MNRAS, 322, 536
[10] Donato D., Ghisellini G., Tagliaferri G., Fossati G., 2001, A&A, 375, 739
[11] Fossati G., Maraschi L., Celotti A., Comastri A., Ghisellini G., 1998, MNRAS, 299, 433
[12] Abdo A. A., et al., 2009, ApJ, 700, 597
[13] Finke J. D., Razzaque S., Dermer C. D., 2010, ApJ, 712, 238
[14] Narumoto, T., & Totani, T. 2006, ApJ, 643, 81
[15] Abdo A. A., et al., 2010, ApJ, 720, 43
[16] Dunlop J. S., Peacock J. A., 1990, MNRAS, 247, 19
[17] Ghirlanda G., Ghisellini G., Tavecchio F., Foschini L., Bonnoli G., 2011, MNRAS, 413, 852
[18] Ghisellini G., et al., 2010, MNRAS, 405, 387
[19] Leroy A. K., Walter F., Brinks E., Bigiel F., de Blok W. J. G., Madore B., Thornley M. D., 2008, AJ, 136, 2782